

MEETING ABSTRACTS

Open Access

# Detection and quantitation of HPV in anogenital and oral tissues and fluids of HIV-positive individuals by real-time PCR

William T Seaman<sup>1\*</sup>, Elizabeth Andrews<sup>2,4</sup>, Marion Couch<sup>1</sup>, Erna Milu Kojic<sup>5</sup>, Susan Cu-Uvin<sup>5</sup>, Allison M Deal<sup>4</sup>, Byrd Quinlavin<sup>5</sup>, Julia Seay<sup>5</sup>, Jennifer Webster-Cyriaque<sup>1,2,3</sup>

From 12<sup>th</sup> International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI)  
Bethesda, MD, USA. 26-27 April, 2010

Human papillomaviruses (HPV) remain a serious world health problem due to their association with anogenital and oral cancers and warts. While over 100 HPV types have been identified, only a subset is associated with malignancy. HPV16 and 18 are the most common oncogenic types, while HPV6 and 11 are the most common types responsible for anogenital warts. These four types cause up to 90% of HPV-associated disease. While other quantitative PCR (qPCR) assays can be used to detect oncogenic HPV, there is no single tube assay that distinguishes the most frequent oncogenic types and the most common types found in warts. A qPCR assay was developed that allowed for detection and quantitation of these 4 HPV types. Type-specific primer pairs and Taq-Man probes allowed single tube multiplex reactions to be performed. Each HPV type was detected over a range from 2 H 10<sup>1</sup> to 2 H 10<sup>6</sup> copies/reaction, providing a reliable method of quantitating type-specific HPV. A Sybr Green-based qPCR assay was developed that utilizes degenerate primers targeting the E1 region of all HPVs. These assays were run in parallel with PCR/sequence gold standard on 76 oral cancers from HIV-negative individuals. Cervical and oral washes were collected from 25 HIV-positive women and 90 HIV-positive men, respectively, being screened for anogenital neoplasia. Samples were analyzed using the newly developed assays. Of the 115 samples, 16% were HPV positive. Cervical washes contained HPV types 44, 67, 35, and 68 and oral specimens contained HPV types 16, 11,

32, 6, 55, 73, and 70. These results indicate that these assays can be used to detect and quantitate HPV in clinical samples obtained by noninvasive measures.

#### Acknowledgements

This work was supported in part by NIDCR OHARA 1 U01 AI068636-01. This article has been published as part of *Infectious Agents and Cancer* Volume 5 Supplement 1, 2010: Proceedings of the 12<sup>th</sup> International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI). The full contents of the supplement are available online at <http://www.biomedcentral.com/1750-9378/5?issue=S1>.

#### Author details

<sup>1</sup>Lineberger Cancer Center, University of North Carolina, Chapel Hill, NC, USA. <sup>2</sup>Division of Infectious Disease, University of North Carolina, Chapel Hill, NC, USA. <sup>3</sup>Department of Microbiology and Immunology, University of North Carolina School of Medicine, Chapel Hill, NC, USA. <sup>4</sup>Department of Dental Ecology, University of North Carolina School of Dentistry, Chapel Hill, NC, USA. <sup>5</sup>Division of Biology and Medicine, Brown University, Providence, RI, USA.

Published: 11 October 2010

doi:10.1186/1750-9378-5-S1-A17

Cite this article as: Seaman *et al.*: Detection and quantitation of HPV in anogenital and oral tissues and fluids of HIV-positive individuals by real-time PCR. *Infectious Agents and Cancer* 2010 **5**(Suppl 1):A17.

\*Correspondence: [tseaman@med.unc.edu](mailto:tseaman@med.unc.edu)

<sup>1</sup>Lineberger Cancer Center, University of North Carolina, Chapel Hill, NC, USA  
Full list of author information is available at the end of the article