

POSTER PRESENTATION

Open Access

Kaposi sarcoma herpesvirus (KSHV)-associated lymphomas are associated with markedly elevated serum IL-10, elevated IL-6, IL-17 and circulating KSHV

Thomas S Uldrick^{1*}, Mark N Polizzotto¹, Kathleen Wyvill¹, Karen Aleman¹, Vickie Marshall⁴, Richard F Little¹, Armando Filie², Mark Raffeld², Seth M Steinberg³, Stefania Pittaluga², Denise Whitby⁴, Robert Yarchoan¹

From 13th International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI)

Bethesda, MD, USA. 7-8 November 2011

Background

KSHV, also called human herpesvirus-8 (HHV8), is the etiologic agent of primary effusion lymphoma (PEL) (including extracavitary variant), and large B-cell lymphoma arising in HHV8-associated multicentric Castleman disease (together, KSHV-associated non-Hodgkin lymphoma (KSHV-NHL). Additional KSHV-associated diseases include: Kaposi sarcoma (KS), a form of multicentric Castleman disease (KSHV-MCD), and a proposed KSHV-associated inflammatory cytokine syndrome (KICS). Like KSHV-MCD, inflammatory symptoms are common in KSHV-NHL. We compared an array of inflammatory and angiogenic cytokines, chemokines, growth factors, and select clinical laboratory values between HIV-infected patients with KSHV-NHL and other lymphomas (HIV-lymphoma).

Methods

Patients were enrolled in HAMB and/or NIAID protocols. Cases had KSHV-NHL; controls other HIV-lymphoma. Clonality of PEL diagnosed from effusions was confirmed by PCR for immunoglobulin rearrangements. Serum was evaluated by ELISA for IFN- γ , IL-1 β , IL-6, IL-8, IL-10, IL-12p70, TNF- α , IL-17, VEGF-A, (Meso-Scale Discovery, Gathersberg, MD), CXCL1, VEGF-C (R&D Systems, Minneapolis, MN). In patients with KSHV-NHL, peripheral blood mononuclear cell associated KSHV viral load was

measured. Clinical data included demographics, CD4 count, albumin, platelets, hemoglobin, and c-reactive protein (CRP). Comparison of each parameter between patients with KSHV-NHL and other HIV-lymphoma employed an exact form of the Wilcoxon rank-sum test. P-values are 2-sided, with $p \leq 0.01$ considered statistically significant, and $0.01 < p < 0.05$ considered strong trends.

Results

Subjects: 13 KSHV-NHL cases: 12 men, 1 woman. Median age 44, (IQR 39-55). 4 white, 4 Hispanic, 3 African-American, 2 African. PEL (8), extracavitary variant PEL (3), large B-cell lymphoma arising in HHV8-associated MCD (2, both with large effusions), history pathology confirmed KSHV-MCD (4). 28 HIV-associated lymphoma controls: 23 men, 5 women. Median age 38 (IQR 35-46). 17 white, 4 Hispanic, 6 African-American, 1 African. Histologies: primary central nervous system lymphoma (13), diffuse large B-cell lymphoma (DLBCL) (10), Hodgkin disease (1), Burkitt lymphoma (2), plasmablastic lymphoma (1), EBV+ large B-cell lymphoma NOS (1). KSHV-NHL subjects had elevated KSHV viral load, [median 2812 copies/ 10^6 cells (IQR 186-115,789)] and CRP [median 51 mg/L (IQR 45-67)]. Compared to other HIV-lymphomas, patients with KSHV-NHL have higher CD4 counts (median CD4 133 vs. 29 cells/uL, $p=0.002$), hypoalbuminemia (median albumin 1.9 vs. 3.5 mg/dL, $p=0.0034$), and trend towards more severe anemia, thrombocytopenia, and hyponatremia. KSHV-NHL is associated with elevated circulating KSHV, marked elevations in IL-10 (513 vs. 12.2 pg/mL, $p < 0.0001$), elevations in IL-6 (29 vs. 4.1 pg/mL,

* Correspondence: uldricks@mail.nih.gov

¹HIV and AIDS Malignancy Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA
Full list of author information is available at the end of the article

$p=0.0013$), IL-17 (1.6 vs. 0.5 pg/mL, $p=0.0074$), and trends towards increased IFN- γ and IL-1 β .

Conclusions

Inflammatory cytokines are important in KSHV-NHL pathogenesis and symptomatology. Clinical and translational studies evaluating these abnormalities in KSHV-associated malignancies are ongoing.

Author details

¹HIV and AIDS Malignancy Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA.

²Laboratory of Pathology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA. ³Biostatistics and Data Management Section, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA. ⁴AIDS and Cancer Virus Program, SAIC-Frederick, National Cancer Institute, National Institutes of Health, Frederick, MD, USA.

Published: 19 April 2012

doi:10.1186/1750-9378-7-S1-P39

Cite this article as: Uldrick *et al.*: Kaposi sarcoma herpesvirus (KSHV)-associated lymphomas are associated with markedly elevated serum IL-10, elevated IL-6, IL-17 and circulating KSHV. *Infectious Agents and Cancer* 2012 **7**(Suppl 1):P39.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

